arXiv:0812.4634v3 [cond-mat.str-el] 3 Apr 2009

Quantum phase transitions in a spin-1/2 alternating Heisenberg antiferromagnetic chain under a staggered transverse magnetic field

Guang-Qiang Zhong, Shou-Shu Gong, Qing-Rong Zheng, Gang Su*
College of Physical Sciences, Graduate University of Chinese Academy of Sciences, P. O. Box 4588, Beijing 100049, China

The magnetic behaviors of a spin-1/2 alternating Heisenberg antiferromagnetic chain in a staggered transverse magnetic field is studied by means of the density-matrix renormalization group method and Jordan-Wigner transformation. Quantum phase transitions of different types are observed in the S=1 Néel and XY-like gapless phases, which result from the competitions between the staggered transverse field and magnetic orders induced by anisotropy and alternating interactions. The results are compared with the mean-field and some exactly resolved results.

PACS numbers: 75.10.Jm, 75.40.Cx

One-dimensional quantum magnets have been at the center of theoretical and experimental attention due to the exotic magnetic properties in condensed matter physics. In particular, the magnetic behaviors of low dimensional quantum magnets in an external magnetic field exhibit many novel characteristics [1]. The gapped Haldane chain [2] compound $\mathrm{Ni(C_2H_8N_2)_2NO_2(ClO_4)}$ in a uniform magnetic field has been widely studied both experimentally [3, 4] and theoretically [5], which shows a commensurate-incommensurate transition as the gap is closed by the field [6]. A S=1 Haldane chain compound $\mathrm{R_2BaNiO_5}$ (R=magnetic rare earth) [7, 8] with an effective staggered magnetic field has been explored theoretically [9, 10] and numerically [11], revealing that the magnetic behavior in a staggered field is totally different from that in a uniform field. Recently, the magnetic properties of a S=1/2 antiferromagnetic (AF)-ferromagnetic (FM) spin chain are extensively studied [12, 13, 14]. This spin chain with nearly the same AF and FM interaction strength has been realized in experiment by the compound $\mathrm{(CH_3)_2NH_2CuCl_3}$ [15]. Hida [16] pointed out that this chain can map onto the S=1 Haldane chain when the FM couplings dominate. Yamanaka *et al.* [17] suggested a phase diagram for the system with an AF anisotropy. The system has the Haldane, S=1 Néel, and XY-like gapless phases for different anisotropies and alternations. In the S=1 Néel and XY-like phases, the gap vanishes and some magnetic orders emerge. In this paper, we shall concentrate primarily on the magnetic properties of the system under a staggered transverse field in various phases. It is expected that the competitions between different factors would yield rich results.

Let us consider a spin-1/2 alternating Heisenberg chain with anisotropy in a transverse staggered magnetic field, as depicted in the inset of Fig. 1. The Hamiltonian reads

$$H = \sum_{j=1}^{N}(S^x_{2j-1}S^x_{2j} + S^y_{2j-1}S^y_{2j} + \lambda S^z_{2j-1}S^z_{2j}) + \beta\sum_{j=1}^{N}\vec{S}_{2j}\cdot\vec{S}_{2j+1} - h_s\sum_{j=1}^{2N}(-1)^j S^x_j, \qquad (1)$$



FIG. 1: (color online) Phase diagram of the spin-1/2 AF-FM alternating quantum spin chain with Hamiltonian (1) [17]. The staggered magnetic behaviors along the arrows will be studied in the context.

where λ stands for the AF anisotropy, $\beta < 0$ (> 0) is the FM (AF) coupling, N is the number of unit cells, and the last term of Eq. (1) introduces the staggered transverse magnetic field. We take the AF coupling as the energy scale and $g\mu_B$=1. The transverse staggered magnetization and susceptibility are defined, respectively, as

$$m_{stag} = \frac{1}{2N}\sum_{j=1}^{2N}(-1)^j\langle S^x_j\rangle, \qquad (2)$$

$$\chi_{stag} = \frac{\partial m_{stag}}{\partial h_s}. \qquad (3)$$

The density-matrix renormalization group (DMRG) method [18, 19] is invoked to study this system. In our calculations, the length of the chain is taken as 60, and the number of the optimal states is kept as 60. We adopt open boundary conditions. The truncation error is less than 10^{-3} in all calculations. The system has an U(1) symmetry in absence of magnetic field, however, the staggered transverse field breaks this symmetry. Thus, there is no good quantum number that can be used to reduce the Hilbert space dimensions in our calculations.

In order to investigate the effects of the anisotropy and FM interaction on m_{stag}, we first calculate exactly m_{stag}

of two S=1/2 spins coupled by XXZ AF and isotropic FM interactions, respectively. For the spins coupled by XXZ AF coupling, m_{stag} behaves as

$$m_{stag}^{AF} = \frac{2h_s}{\sqrt{16h_s^2 + (1+\lambda)^2}}. \tag{4}$$

It can be seen that with increasing λ, m_{stag} is supressed by the anisotropy that destructs the transverse magnetic ordering. For the spins coupled by the FM interaction, m_{stag} behaves as

$$m_{stag}^{FM} = \frac{h_s}{\sqrt{4h_s^2 + \beta^2}}. \tag{5}$$

Clearly, the FM interaction makes the spins align in the same direction, which competes with the staggered magnetic field. Thus, m_{stag} is supressed with increasing β. It seems that the anisotropy and FM interaction have similar effects on m_{stag}. However, when the FM coupling and the staggered magnetic field are considered simultaneously, m_{stag} shows more complex behaviors. In the following, the transverse staggered magnetic properties will be studied by means of the DMRG method for the parameters indicated by the arrows of Fig. 1.

Figure 2 shows the evolution of the magnetic properties from the Haldane phase to the S=1 Néel phase with changing β from β=1 to β=−4 for λ=4.0. In Fig. 2(a), m_{stag} increases with h_s and approaches 0.5 when $h_s \to \infty$. With increasing FM interactions, m_{stag} declines for any field, which agrees with the depressing effect of the FM coupling. In the Haldane phase, χ_{stag} continuously declines with h_s in a simple way

$$\chi_{stag} \sim \frac{1}{(h_s^2 + A)^{3/2}}, \tag{6}$$

where A is a constant determined by the couplings. At h_s=0, the zero-field susceptibility $\chi_{stag}(0)$ is finite, which is analogous to the S=1 Haldane chain [7, 8, 9, 11] due to the existence of a gap. For the S=1 Haldane chain, $\chi_{stag}(0) = Zv/\Delta^2$ (Δ is the spin gap) [9]. In the present case, the spin gap [16] and $\chi_{stag}(0)$ decrease simultaneously with increasing FM coupling. Therefore, it is expected that the product of the renormalization parameter Z and the spin-wave velocity v decrease more rapidly than the gap. With further increasing the FM coupling, χ_{stag} becomes flat at low fields when approaching the phase boundary. In the S=1 Néel phase, a broad peak emerges in χ_{stag} [Fig. 2(b)], indicating the distinct magnetic properties below the transition field from those in the Haldane phase. The peaks, which move to higher fields with increasing $-\beta$, is attributed to the competition between the field and magnetic interactions. In the absence of magnetic field, the longitudinal spin-spin correlation function $\langle S_0^z S_j^z \rangle$ has a long-range order (LRO) and behaves as $\langle S_0^z S_{2j-1}^z \rangle \simeq \langle S_0^z S_{2j}^z \rangle$ due to the FM coupling [Fig. 2(c)]. The Néel order in the z axis prevents the magnetization in the x direction and is destructed by the transverse staggered field. As shown in Fig. 2(c), with increasing h_s, $\langle S_0^z S_j^z \rangle$ decays with a power law, and when h_s exceeds a critical magnetic field, h_{s_c}, it decays exponentially. Below h_{s_c}, the suppressed Néel order facilitates the magnetization in the transverse axis, yielding the increase of χ_{stag}. Above h_{s_c}, the Néel order is fully broken and thus χ_{stag} declines similar to that in the Haldane phase. In the Haldane phase, the gap enlarges with increasing h_s. In the Néel phase, a gap is opened when h_s exceeds h_{s_c} and increases with the field, as shown in Fig. 2(d). In the transition field h_{s_c}, a quantum phase transition (QPT) [20] which is from the S=1 Néel phase to a gapped staggered magnetic ordered phase happens. In the vicinity of h_{s_c}, the ground-state energy e and its derivatives with respect to the field are studied to characterize this transition. It is found that both $\partial e/\partial h_s$ and $\partial^2 e/\partial {h_s}^2$ are continuous and nonsingular in the field h_{s_c}, but $\partial^2 e/\partial {h_s}^2$ has a minimum at h_{s_c}, as shown in the inset of Fig. 2(d). Below h_{s_c}, the phase is gapless with a power law decaying $\langle S_0^z S_j^z \rangle$. Above h_{s_c}, a gap emerges and $\langle S_0^z S_j^z \rangle$ decays exponentially. This is analogous to the transition from the XY-like phase to the Haldane phase in the absence of the field [21], which is of the Kosterlitz-Thouless (KT) [22] type. By considering the nonsingularity of the derivatives of e, we argue that this QPT may be also of the KT type. The critical behavior of Δ near h_{s_c} is fitted by the KT type with the lines in Fig. 2(d)

$$\Delta = De^{-C/\sqrt{h_s - h_{s_c}}}. \tag{7}$$

For β=−3(−4), D=2.7(3.4), C=2.0(2.0), h_{s_c}=0.7(1.27). The behavior of Δ near h_{s_c} can be fitted well by Eq. (7).

Figure 3 shows the changes of magnetic properties from the Haldane phase to the XY-like gapless phase with changing β from β=1 to β=−5 for λ=−0.5. In the XY-like gapless phase, m_{stag} has an inflexion with increasing h_s, which is explicitly characterized by the sharp peak of χ_{stag} and indicates a transition of magnetic properties. Like the Néel phase, the peak moves to higher fields with increasing $-\beta$ [Fig. 3(b)]. This transition is due to the competition between the magnetic field and transverse magnetic ordering. In the absence of magnetic field, $\langle S_0^x S_j^x \rangle$ has a quasi-LRO, and due to the FM interaction, $\langle S_0^x S_{2j-1}^x \rangle \simeq \langle S_0^x S_{2j}^x \rangle$, i.e., $\langle S_0^x S_j^x \rangle$ has a translation symmetry with a period of 4 [Fig. 3(c)]. As the staggered field competes with the FM interaction, this transverse quasi-LRO prevents the staggered magnetization, and meanwhile, is destructed by increasing the field. With increasing h_s, $\langle S_0^x S_j^x \rangle$ becomes disordered in short range but builds an order with a 2-period translation symmetry in long range, as shown in Fig. 3(c) for h_s=0.5. When h_s exceeds the transition field, the short-range disorder is replaced by the staggered magnetic ordering, as in the case of h_s=1.4 in Fig. 3(c). These variations are also visible in



FIG. 2: (color online) Field dependence of m_{stag}, χ_{stag}, $\langle S_0^z S_j^z\rangle$, and Δ for λ=4.0 in (a), (b), (c), and (d), respectively. The lines in (d) denote the fitted results with Eq. (7). The inset of (d) shows the second derivative of the ground-state energy $\partial^2 e/\partial {h_s}^2$ as a function of h_s.

spin static structure factor $S(q)$, which are not presented here for concise. In the absence of magnetic field, $S(q)$ has two peaks at q=$\pi/2$ and $3\pi/2$. With increasing h_s, the old peaks decline rapidly and a new peak emerges at q=π, indicating the changing periodicity of $\langle S_0^x S_j^x\rangle$. Below h_{s_c}, the destructed quasi-LRO facilitates the staggered magnetization and thus χ_{stag} increases. Above h_{s_c}, the quasi-LRO is fully broken and χ_{stag} declines as in the Haldane phase. Similar to the S=1 Néel phase, this transition is also accompanied by the open of a gap, which, however, behaves in a different manner above h_{s_c}

$$\Delta \sim (h_s - h_{s_c})^{\alpha}. \tag{8}$$

The critical behaviors of Δ are fitted by Eq. (8) in Fig. 3(d) with α=1.15 and 1.1 for β=-3 and -5, respectively. These observations indicate that a QPT happens in h_{s_c}. To characterize this QPT, the ground-state energy and its derivatives with the field are studied. It is found that both e and $\partial e/\partial h_s$ are nonsingular, but $\partial^2 e/\partial {h_s}^2$ is singular at h_{s_c}, indicating that this transition is of the second-order, as shown in the inset of Fig. 3(d).

As discussed above, the staggered transverse magnetic properties have different behaviors in three phases. Although the QPT is observed in the S=1 Néel and XY-like phases, the physical quantities have different behaviors in these phases. The QPT in the Néel phase is argued to be of the KT type, while that in the XY-like phase is confirmed to be of the second-order. For further discussions, the staggered magnetization is studied in terms of the Jordan-Wigner (JW) transformation. As the field is applied transversely, the transformation is introduced as



FIG. 3: (color online) Field dependence of m_{stag}, χ_{stag}, $\langle S_0^x S_j^x\rangle$, and Δ for β=-0.5 in (a), (b), (c), and (d), respectively. The inset of (d) shows $\partial^2 e/\partial {h_s}^2$ as a function of h_s.

$$\begin{aligned} S_i^z &= \frac{1}{2}(c_i^{\dagger} e^{i\pi\sum_{j<i} c_j^{\dagger}c_j} + h.c.), \\ S_i^y &= \frac{1}{2i}(e^{-i\pi\sum_{j<i} c_j^{\dagger}c_j} c_i - h.c.), \\ S_i^x &= c_i^{\dagger}c_i - \frac{1}{2}, \end{aligned} \tag{9}$$

where $c_i^{\dagger}$ and c_i are the creation and annihilation operators of spinless fermions, which satisfy the anticommutation relation $\{c_i, c_j^{\dagger}\}$=δ_{ij}. We denote the fermions in odd and even sites as a_i and b_i, respectively. The density-density interaction terms are treated with the Hartree-Fock (HF) approximation,

$$\begin{aligned} n_{a,j}n_{b,j} &\approx n_{a,j}\langle n_{b,j}\rangle + \langle n_{a,j}\rangle n_{b,j} - (a_j^{\dagger}b_j\langle b_j^{\dagger}a_j\rangle + h.c.) \\ &+ (a_j^{\dagger}b_j^{\dagger}\langle b_j a_j\rangle + h.c.) - (\langle n_{a,j}\rangle\langle n_{b,j}\rangle \\ &- \langle b_j^{\dagger}a_j\rangle\langle a_j^{\dagger}b_j\rangle + \langle b_j a_j\rangle\langle a_j^{\dagger}b_j^{\dagger}\rangle). \end{aligned} \tag{10}$$

We denote $\langle a_i^{\dagger}a_i\rangle$=$n_a$, $\langle b_i^{\dagger}b_i\rangle$=$n_b$, $\langle b_i a_i\rangle$=p_1, $\langle b_i^{\dagger}a_i\rangle$=$p_2$, $\langle a_{i+1}^{\dagger}b_i\rangle$=$p_3$, and $\langle a_{i+1}b_i\rangle$=$p_4$. After making Fourier transform, the Hamiltonian is transformed into

$$\begin{aligned} H_{HF} &= \sum_k(\omega_k^a a_k^{\dagger}a_k + \omega_k^b b_k^{\dagger}b_k) + \sum_k(\omega_k^1 a_k^{\dagger}b_k + \omega_k^2 a_k^{\dagger}b_{-k}^{\dagger} \\ &+ h.c.) + const., \end{aligned} \tag{11}$$

where ω_k^a=$(n_b - 1/2)(\beta+1) - h_s$, ω_k^b=$(n_a - 1/2)(\beta + 1) + h_s$, ω_k^1=$[(\lambda+1)/4 - p_2]e^{ik/2}$+$(1/2 - p_3^*)\beta e^{-ik/2}$, and ω_k^2=$[(\lambda-1)/4 + p_1]e^{ik/2}$-$p_4^*\beta e^{-ik/2}$. Then, we introduce the Bogoliubov transformation

$$a_k = u_{11}\alpha_k + u_{12}\beta_k + u_{13}\gamma_k + u_{14}\lambda_k,$$



FIG. 4: (color online) Staggered magnetization and susceptibility obtained by (a) the mean-field theory for the Hamiltonian Eq. (13); and (b) the exact solutions for the Hamiltonian Eq. (1) with only Ising interactions.

$$\begin{aligned} b_k &= u_{21}\alpha_k + u_{22}\beta_k + u_{23}\gamma_k + u_{24}\lambda_k, \\ a^\dagger_{-k} &= u_{31}\alpha_k + u_{32}\beta_k + u_{33}\gamma_k + u_{34}\lambda_k, \\ b^\dagger_{-k} &= u_{41}\alpha_k + u_{42}\beta_k + u_{43}\gamma_k + u_{44}\lambda_k. \end{aligned} \tag{12}$$

Therefore, the Hamiltonian is diagonalized as

$$H_{HF} = \sum_k (\omega^\alpha_k \alpha^\dagger_k \alpha_k + \omega^\beta_k \beta^\dagger_k \beta_k + \omega^\gamma_k \gamma^\dagger_k \gamma_k + \omega^\lambda_k \lambda^\dagger_k \lambda_k), \tag{13}$$

where ω^i_k (i=$\alpha, \beta, \gamma, \lambda$) are obtained by the self-consistent numerical calculations. The staggered magnetic properties are studied with this quardratic Hamiltonian. In the following, we find that this mean-field theory is able to describe the behavior in the Haldane phase, but fails to reproduce the QPT in other phases. Figure 4(a) shows m_{stag} and χ_{stag} from the Haldane to the Néel phase. The analytic results qualitatively agree with the numerical ones in the Haldane phase, but cannot find the inflexion in the Néel phase. However, some exactly soluble case that has a competition between the field and magnetic couplings also shows QPT and may be compared with our observations. Figure 4(b) shows the exact solutions for the Ising case of Eq. (1), which exhibits a QPT with increasing h_s. The transition behaviors are analogous to those in the XY-like phase. The singular $\partial^2 e/\partial {h_s}^2$ in h_{s_c} indicates that this QPT also belongs to the second-order, which are not presented here. The gap above h_{s_c} behaves as $\Delta \sim (h_s - h_{s_c})$, which is Eq. (8) with α=1. It may be expected that the QPT in the Ising case is similar to that in the S=1 Néel phase, but it is of the second-order, like the transition in the XY-like phase. This phenomenon shows the important role of the xy-component quantum fluctuations in the QPT, which *smooth* the transition in the Néel phase.

In summary, we have studied the magnetic properties of the spin-1/2 AF-FM Heisenberg chain with AF anisotropy in a transverse staggered magnetic field by means of the DMRG method. The physical quantities are explored in the Haldane, S=1 Néel, and XY-like gapless phases. In the Haldane phase, m_{stag} and χ_{stag} behave as those of the S=1 Haldane chain, and do not have transition behaviors. In the Néel phase and the XY-like gapless phase, due to the competitions between the field and different magnetic couplings, most quantities have a transition at a field h_{s_c}, indicating a QPT induced by the field happens in the system. m_{stag} has an inflexion in h_{s_c}, which corresponds to a maximum in χ_{stag}. The transition is also accompanied by the open of a gap. In the Néel phase, $\partial^2 e/\partial {h_s}^2$ is nonsingular, but it still describes the transition with a minimum at the critical field. The QPT is argued to be of the KT type. The critical behavior of Δ is well fitted by that in the KT transition near h_{s_c}. In the XY-like gapless phase, $\partial^2 e/\partial {h_s}^2$ is singular in h_{s_c}, indicating that the transition is of the second order. The gap near h_{s_c} behaves as $\Delta \sim (h_s - h_{s_c})^\alpha$. Due to the distinct magnetic orders, the transitions behave differently in the two phases. Using the Jordan-Wigner transformation, the magnetic properties are also investigated analytically. In the present HF approximation, the features in the Haldane phase are reproduced, but it fails to describe the transitions in other two phases. The Ising case of the Hamiltonian (1) is exactly resolved, which has a second-order QPT that behaves like the one in the XY-like phase. The differences between the Ising case and the Néel phase show the important role of the quantum fluctuations.

We are grateful to Wei Li and Yang Zhao for useful discussions. This work is supported in part by the National Science Fund for Distinguished Young Scholars of China (Grant No. 10625419), the National Science Foundation of China (Grants No. 90403036 and No. 20490210), the MOST of China (Grant No. 2006CB601102), and the Chinese Academy of Sciences.

[] *Corresponding author. E-mail: gsu@gucas.ac.cn

[1] For instance, B. Gu, G. Su, S. Gao, J. Phys.:Condens. Matter **17**, 6081 (2005); B. Gu, G. Su, S. Gao, Phys. Rev. B **73**, 134427 (2006); B. Gu and G. Su, Phys. Rev. Lett. **97**, 089701 (2006); B. Gu and G. Su, Phys. Rev. B **75**, 174437 (2007); S. S. Gong, B. Gu, G. Su, Phys. Lett. A **372**, 2322 (2008); S. S. Gong and G. Su, Phys. Rev. B **78**, 104416 (2008).

[2] F. D. Haldane, Phys. Rev. Lett. **50**, 1153 (1983); Phys. Lett. **93A**, 464 (1983).

[3] K. Katsumata *et al.*, Phys. Rev. Lett. **63**, 86 (1989).

[4] M. Chiba *et al.*, Phys. Rev. B **44**, 2838 (1991).

[5] I. Affleck, Phys. Rev. B **41**, 6697 (1990).

[6] R. Chitra and G. Giamarchi, Phys. Rev. B **55**, 5816 (1997).

[7] A. Zheludev, E. Ressouche, S. Maslov, T. Yokoo, S. Raymond, and J. Akimitsu, Phys. Rev. Lett. **80**, 3630 (1998).

[8] S. Maslov and A. Zheludev, Phys. Rev. B **57**, 68 (1998).

[9] S. Maslov and A. Zheludev, Phys. Rev. Lett. **80**, 5786 (1998).

[10] E. Ercolessi, G. Morandi, P. Pieri, and M. Roncaglia, Phys. Rev. B **62**, 14860 (2000).

[11] J. Z. Lou, X. Dai, S. J. Qin, Z. B. Su, and L. Yu, Phys. Rev. B **60**, 52 (1999).

[12] T. Sakai, J. Phys. Soc. Jpn. **64**, 251 (1995).

[13] S. Mahdavifar and A. Akbari, J. Phys. Soc. Jpn. **77**, 024710 (2008).

[14] J. Abouie and S. Mahdavifar, Phys. Rev. B **78**, 184437 (2008).

[15] M. B. Stone *et al.*, Phys. Rev. Lett. **99**, 087204 (2007).

[16] K. Hida, Phys. Rev. B **45**, 2207 (1992).

[17] M. Yamanaka, Y. Hatsugai, and M. Kohmoto, Phys. Rev. B **48**, 9555 (1993).

[18] S. R. White, Phys. Rev. Lett. **69**, 2863 (1992); S. R. White, Phys. Rev. B **48**, 10345 (1993).

[19] U. Schollwöck, Rev. Mod. Phys. **77**, 259 (2005).

[20] S. Sachdev, *Quantum Phase Transitions* (Cambridge University Press, Cambridge, 1999).

[21] A. Kitazawa, K. Nomura, and K. Okamoto, Phys. Rev. Lett. **76**, 4038 (1996).

[22] J. M. Kosterlitz and D. J. Thouless, J. Phys. C **6**, 1181 (1973).